arXiv:math/0301105v2 [math.DG] 11 Jan 2003

Rigid 6-dimensional h-spaces of constant curvature[1]

Zolfira Zakirova[2]

Abstract

In this paper, we continue studying the 6-dimensional pseudo-Riemannian space $V^6(g_{ij})$ with signature $[++----]$, which admits projective motions, i. e. continuous transformation groups preserving geodesics. In particular, we determine a necessary and sufficient condition that the 6-dimensional rigid h-spaces have constant curvature.

1 Introduction

The general method of determining pseudo-Riemannian manifolds that admit some nonhomothetic projective group G_r has been developed by A.V.Aminova [A]. A.V.Aminova has classified all the Lorentzian manifolds of dimension ≥ 3 that admit nonhomothetic projective or affine infinitesimal transformations. In each case, there were determined the corresponding maximal projective and affine Lie algebras. This problem is not solved for pseudo-Riemannian spaces with arbitrary signature. In the series of works, we investigate the 6-dimensional pseudo-Riemannian space $V^6(g_{ij})$ with signature $[++----]$.

Remind that in order to find a pseudo-Riemannian space admitting a nonhomothetic infinitesimal projective transformation, one needs to integrate the Eisenhart equation [E]

$$h_{ij,k} = 2g_{ij}\varphi_{,k} + g_{ik}\varphi_{,j} + g_{jk}\varphi_{,i}. \tag{1}$$

Pseudo-Riemannian manifolds for which there exist nontrivial solutions $h_{ij} \neq cg_{ij}$ to the Eisenhart equation are called *h-spaces.* It is known that the problem of determining such spaces depends on the type of the h-space, i. e. on the type of the bilinear form $L_X g_{ij}$ determined by the characteristic of the λ-matrix $(h_{ij} - \lambda g_{ij})$. The number of possible types

---

[1] The research was partially supported by the RFBR grant 01-02-17682-a and by the INTAS grant 00-00334.

[2] Kazan State University, e-mail: Zolfira.Zakirova@soros.ksu.ru, Zolfira.Zakirova@ksu.ru

depends on the dimension and the signature of the h-space. In our case of six-dimensional spaces V^6 with signature $[++----]$, there is a variety of possible types. We restrict it considering the rigid h-spaces: h-spaces with distinct bases of prime divisors of the λ-matrix are called rigid h-spaces [P, Z]. In the papers [Z1]-[Z4] (see also [Z5]), we found 6-dimensional rigid h-spaces of the [2211], [321], [33], [411] and [51] types.

For further study we need a necessary and sufficient condition of constant curvature of this h-space. This condition is known to be expressed by the formula

$$R^i_{jkl} = K(\delta_k{}^i g_{jl} - \delta_l{}^i g_{jk}), \quad K = \text{const}. \tag{2}$$

Since the projective-group properties of the rigid h-spaces of the [21111] and [3111] types have been investigated by A. V. Aminova, we do not discuss them here.

2 The h-space of the $[2211]$ type

Let us investigate the h-space of the [2211] type. In this case, metric has the following form

$$g_{ij}dx^i dx^j = e_2(f_4 - f_2)^2 \Pi_\sigma (f_\sigma - f_2)\{2A dx^1 dx^2 - A^2 \Sigma_1 (dx^2)^2\} + \tag{3}$$

$$e_4(f_2 - f_4)^2 \Pi_\sigma (f_\sigma - f_4)\{2\tilde{A} dx^3 dx^4 - \tilde{A}^2 \Sigma_2 (dx^4)^2\} + \sum_\sigma e_\sigma \Pi'_i (f_i - f_\sigma)(dx^\sigma)^2,$$

where

$$A = \epsilon x^1 + \theta(x^2), \quad \tilde{A} = \tilde{\epsilon} x^3 + \omega(x^4), \tag{4}$$

$$\Sigma_1 = 2(f_4 - f_2)^{-1} + \sum_\sigma (f_\sigma - f_2)^{-1}, \quad \Sigma_2 = 2(f_2 - f_4)^{-1} + \sum_\sigma (f_\sigma - f_4)^{-1},$$

$f_2 = \epsilon x^2$, $f_4 = \tilde{\epsilon} x^4 + a$, $\epsilon, \tilde{\epsilon} = 0, 1$, a is a constant nonzero when $\tilde{\epsilon} = 0$, $f_\sigma(x^\sigma)$, $\theta(x^2)$, $\omega(x^4)$ are arbitrary functions, $e_i = \pm 1$.

Calculating components of the curvature tensor of our h-space, we observe that the nonzero components are

$$R^{a_p}_{b_p a_p c_p} = \chi_p g_{b_p c_p}, \tag{5}$$

$$R^{\sigma}_{b_p \sigma c_p} = \rho_{\sigma p} g_{a_p b_p} - \frac{\chi_p - \rho_{\sigma p}}{f_\sigma - f_p} A_p g_{a_p c_p} \delta^p_{b_p}, \ (c_p \neq b_p),$$

$$R^{a_p}_{\sigma b_p \sigma} = g_{\sigma\sigma}\{\rho_{\sigma p} \delta^{a_p}_{b_p} - \frac{\chi_p - \rho_{\sigma p}}{f_\sigma - f_p} A_p \delta^p_{b_p} \delta^{a_p}_{c_p}\}, \ (c_p \neq b_p),$$

$$R^{\tau}_{\sigma\tau\sigma} = g_{\sigma\sigma} \frac{\rho_{\tau p}(f_\tau - f_p) - \rho_{\sigma p}(f_\sigma - f_p)}{f_\tau - f_\sigma}, \ (\sigma \neq \tau),$$

$$R^{a_p}_{a_q b_p b_q} = \{\rho_{pq} g_{a_q b_q} - \frac{\chi_q - \rho_{pq}}{f_p - f_q} A_q g_{a_q c_q} \delta^q_{b_q}\} \delta^{a_p}_{b_p} -$$

$$\{\frac{\chi_p - \rho_{pq}}{f_q - f_p} A_p g_{a_q b_q} + \frac{\sum\limits_{l=2,4} (\chi_l - \rho_{pq})}{(f_q - f_p)^2} A_p A_q g_{a_q c_q} \delta^q_{b_q}\} \delta^p_{b_p} \delta^{a_p}_{c_p},$$

$$(c_p \neq b_p, c_q \neq b_q, p \neq q),$$

where

$$\chi_p = B_p + \rho_p, \quad B_2 = \frac{\epsilon\theta'}{A^2 g_{12}}, \; B_4 = \frac{\tilde{\epsilon}\omega'}{\tilde{A}^2 g_{34}}, \quad A_2 = A, \; A_4 = \tilde{A}, \tag{6}$$

$$\rho_p = -\frac{1}{4}\sum_{\sigma} \frac{(f'_\sigma)^2}{(f_\sigma - f_p)^2 g_{\sigma\sigma}}, \quad \rho_{pq} = -\frac{1}{4}\sum_{\sigma} \frac{(f'_\sigma)^2}{(f_\sigma - f_p)(f_\sigma - f_q) g_{\sigma\sigma}}, \tag{7}$$

$$\rho_{\sigma p} = -\frac{1}{4}\frac{(f'_\sigma)^2}{(f_\sigma - f_p) g_{\sigma\sigma}} \{\frac{2f''{}_\sigma}{(f'_\sigma)^2} - \frac{1}{f_\sigma - f_p} + \sum_{i, i \neq \sigma} (f_i - f_\sigma)^{-1}\} -$$

$$\frac{1}{4}\sum_{\gamma, \gamma \neq \sigma} \frac{(f'_\gamma)^2}{(f_\gamma - f_p)(f_\gamma - f_\sigma) g_{\gamma\gamma}}.$$

Here $p, q = 2, 4$, $\sigma = 5, 6$, the indices a_2 are equal to either 1 or 2, while the indices a_4 are 3 or 4.

In particular, from (2) it follows that

$$R^1_{112} = R^\sigma_{1\sigma 2}, \quad R^3_{334} = R^\sigma_{3\sigma 4}, \quad R^1_{112} = R^4_{142}, \quad R^3_{334} = R^2_{324}.$$

Substituting components of the curvature tensor into this equality, one obtains

$$(\chi_2 - \rho_{\sigma 2}) g_{12} = 0, \quad (\chi_4 - \rho_{\sigma 4}) g_{34} = 0, \quad (\chi_2 - \rho_{24}) g_{12} = 0, \quad (\chi_4 - \rho_{24}) g_{34} = 0,$$

since $g_{12} \neq 0$ and $g_{34} \neq 0$, then

$$\chi_2 - \rho_{\sigma 2} = 0, \quad \chi_4 - \rho_{\sigma 4} = 0, \tag{8}$$

$$\chi_2 - \rho_{24} = 0, \quad \chi_4 - \rho_{24} = 0.$$

Let us prove that the condition $\chi_2 - \rho_{\sigma 2} = 0$ is equivalent to the conditions

$$\rho_2 - \rho_{\sigma 2} = \epsilon = 0.$$

Indeed, differentiating the equality

$$\chi_2 - \rho_{\sigma 2} = \rho_2 - \rho_{\sigma 2} + \frac{\epsilon\theta'}{A^2 g_{12}} = 0$$

with respect to x^1, we obtain $\epsilon = 0$ and, therefore, $\rho_2 - \rho_{\sigma 2} = 0$. Inversely, if $\rho_2 - \rho_{\sigma 2} = \epsilon = 0$, then $\chi_2 - \rho_{\sigma 2} = 0$. Similarly we can find equivalent conditions for other relations (8).

Let us prove that the necessary conditions of constant curvature of the h-space of the [2211] type,

$$\chi_p - \rho_{\sigma p} = \chi_p - \rho_{24} = 0, \tag{9}$$

or equivalent conditions

$$\rho_p - \rho_{\sigma p} = \rho_p - \rho_{24} = \epsilon = \tilde{\epsilon} = 0, \tag{10}$$

are sufficient. The relations that follows from (7),

$$\partial_\sigma \rho_p = -f'_\sigma \frac{\rho_p - \rho_{\sigma p}}{f_\sigma - f_p}, \quad \partial_1 \rho_p = \partial_3 \rho_p = 0,$$

$$\partial_p \rho_p = -\epsilon_p \sum_\sigma \frac{{f'_\sigma}^2}{(f_\sigma - f_p)^3 g_{\sigma\sigma}}, \quad \partial_p \rho_q = -\frac{\epsilon_p}{2} \sum_\sigma \frac{{f'_\sigma}^2}{(f_\sigma - f_q)^2 (f_\sigma - f_p) g_{\sigma\sigma}},$$

where $p \neq q$, $\epsilon_2 = \epsilon$, $\epsilon_4 = \tilde{\epsilon}$, along with (10), lead to the identities

$$\chi_p = \rho_{\sigma p} = \rho_{24} = \rho_p = \text{const.}$$

Using these results along with conditions $\epsilon = \tilde{\epsilon} = 0$, one can prove that components of the curvature tensor (5) satisfy conditions (2) with $K = \rho_p = \text{const.}$

3 The h-space of the [321] type

Metric of this h-space can be written in the form

$$g_{ij} dx^i dx^j = e_3 (f_5 - \epsilon x^3)^2 (f_6 - \epsilon x^3) \{ (dx^2)^2 + 4A dx^1 dx^3 + \tag{11}$$

$$2(\epsilon x^1 - 2A\Sigma_1) dx^2 dx^3 + ((\epsilon x^1)^2 - 4\epsilon x^1 A \Sigma_1 + 4A^2 \Sigma_3)(dx^3)^2 \} +$$

$$e_5 (\epsilon x^3 - f_5)^3 (f_6 - f_5) \{ 2\tilde{A} dx^4 dx^5 - \Sigma_4 \tilde{A}^2 (dx^5)^2 \} + e_6 (f_5 - f_6)^2 (f_5 - f_6)^3 (dx^6)^2,$$

where

$$A = \epsilon x^2 + \theta(x^3), \quad \tilde{A} = \tilde{\epsilon} x^4 + \omega(x^5), \tag{12}$$

$$\Sigma_1 = (f_6 - \epsilon x^3)^{-1} + 2(f_5 - \epsilon x^3)^{-1}, \quad \Sigma_2 = (f_6 - \epsilon x^3)^{-2} + 2(f_5 - \epsilon x^3)^{-2},$$

$$2\Sigma_3 = (\Sigma_1)^2 - \Sigma_2, \quad \Sigma_4 = 3(\epsilon x^3 - f_5)^{-1} + (f_6 - f_5)^{-1},$$

$f_1 = f_2 = f_3 = \epsilon x^3$, $f_4 = f_5 = \tilde{\epsilon} x^5 + a$, f_6 is an arbitrary function of the variable x^6, $\epsilon, \tilde{\epsilon} = 0, 1$, $\epsilon \neq 0$ when $\tilde{\epsilon} = 0$, and, inversely, $\tilde{\epsilon} \neq 0$ when $\epsilon = 0$, a is a constant nonequal to zero when $\tilde{\epsilon} = 0$, $e_3, e_5, e_6 = \pm 1$.

From (2) one finds

$$R^2_{123} = R^r_{1r3}, \quad (r \;=\; 5,6).$$

Substituting into this equality the corresponding components of the curvature tensor of our h-space

$$R^2_{123} = \chi_3 g_{13}, \quad R^5_{153} = \rho_{53} g_{13}, \quad R^6_{163} = \rho_{63} g_{13},$$

where

$$\chi_3 = \frac{3\epsilon^2}{16A^2 g_{22}} + \rho_3, \tag{13}$$

ρ_3, ρ_{53} and ρ_{63} are defined by formulas (7) with $p,q = 3,5$ and $\sigma = 5,6$, one gets

$$(\chi_3 - \rho_{r3}) g_{13} = 0.$$

Since $g_{13} \neq 0$, then

$$\chi_3 - \rho_{r3} = \frac{3\epsilon^2}{16A^2 g_{22}} + \rho_3 - \rho_{r3} = 0.$$

Differentiating the last equality with respect to x^2, one obtains $\epsilon = 0$ and, therefore, $\rho_3 - \rho_{r3} = 0$. Inversely, if $\rho_3 - \rho_{r3} = \epsilon = 0$, then $\chi_3 - \rho_{r3} = 0$. Since $f_3 \neq f_5$, from $\rho_3 - \rho_{53} = 0$ it follows that $f'_6 = 0$. Substituting $\epsilon = 0$ into the relation

$$R^1_{123} = \gamma g_{13} + \frac{3\epsilon^2}{8A}\Sigma_1 + \frac{3\epsilon}{4A^2}(\theta' - \epsilon^2 x^1) = 0,$$

where $R^1_{123} = 0$ follows from (2), one finds

$$\gamma = -\frac{1}{4}\sum_\sigma \frac{(f'_\sigma)^2}{(f_\sigma - f_p)^3 g_{\sigma\sigma}} = 0.$$

From (2), one has

$$R^4_{445} = R^s_{4s5}, \quad (s = 3,6).$$

Therefore,

$$(\chi_5 - \rho_{s5}) g_{45} = 0,$$

since for our h-space

$$R^4_{445} = \chi_5 g_{45}, \quad R^3_{435} = \rho_{53} g_{45}, \quad R^6_{465} = \rho_{65} g_{45},$$

where

$$\chi_5 = \frac{\tilde{\epsilon}\theta'}{\tilde{A}^2 g_{45}} + \rho_5. \tag{14}$$

Since $g_{45} \neq 0$, we obtain $\chi_5 - \rho_{s5} = 0$. Similarly one can prove that this condition is equivalent to the conditions $f'_6 = \tilde{\epsilon} = 0$.

Let us prove that the necessary conditions of constant curvature of the h-space of the [321] type,

$$\chi_p - \rho_{6p} = \chi_p - \rho_{35} = \gamma = 0, \quad (p = 3, 5), \tag{15}$$

or

$$f_6' = \epsilon = \tilde{\epsilon} = 0,$$

are sufficient. Note that from (15) and (7) ($p, q = 3, 5, \sigma = 6$), (13) and (14) it follows that

$$\chi_p = \rho_p = \rho_{6p} = \rho_{35} = 0.$$

Calculating the other components of the curvature tensor of our h-space and substituting them into the last equalities, one gets h-space of the constant curvature.

4 The h-space of the [33] type

Metric is determined by formulas

$$g_{ij} dx^i dx^j = e_3 (f_6 - f_3)^3 \{ (dx^2)^2 + 4A dx^1 dx^3 + \tag{16}$$

$$2(\epsilon x^1 - 2A\Sigma_1) dx^2 dx^3 + ((\epsilon x^1)^2 - 4\epsilon x^1 A \Sigma_1 + 4A^2 \Sigma_2)(dx^3)^2 \} +$$

$$e_6 (f_3 - f_6)^3 \{ (dx^5)^2 + 4\tilde{A} dx^4 dx^6 + 2(\tilde{\epsilon} x^4 + 2\tilde{A}\Sigma_1) dx^5 dx^6 + ((\tilde{\epsilon} x^4)^2 + 4\tilde{\epsilon} x^4 \tilde{A} \Sigma_1 + 4\tilde{A}^2 \Sigma_2 ((dx^6)^2 \},$$

where

$$A = \epsilon x^2 + \theta(x^3), \quad \tilde{A} = \tilde{\epsilon} x^4 + \omega(x^6), \tag{17}$$

$$\Sigma_1 = 3(f_6 - f_3)^{-1}, \quad \Sigma_2 = 3(f_6 - f_3)^{-2},$$

$f_3 = \epsilon x^3$, $f_6 = \tilde{\epsilon} x^6 + a$, $\epsilon, \tilde{\epsilon} = 0, 1$, c is a constant, a is a constant nonequal to zero when $\tilde{\epsilon} = 0$, $i_1, j_1 = 1, 2, 3$, $i_2, j_2 = 4, 5, 6$, $e_i = \pm 1$, $\theta(x^3)$, $\omega(x^6)$ are arbitrary functions, nonzero when $\epsilon = 0$ and $\tilde{\epsilon} = 0$ respectively.

Nonzero components $R^{a_p}_{\nu b_p \mu}$ ($b_p \neq \mu, a_p < b_p$), $R^{a_1}_{5a_1 6}$, $R^{a_1}_{6a_1 5}$, $R^{a_1}_{6a_1 6}$, $R^{a_2}_{2a_2 3}$, $R^{a_2}_{3a_2 2}$, $R^{a_2}_{3a_2 3}$ of the curvature tensor ($p = 1, 2$, $a_1, b_1 = 1, 2, 3$, $a_2, b_2 = 4, 5, 6$, ν and μ are the indices of nonzero components of metric $g_{\nu\mu}$ (16)) are proportional to ϵ or to $\tilde{\epsilon}$. In particular,

$$R^2_{123} = \frac{3\epsilon^2}{8A}, R^5_{456} = \frac{3\tilde{\epsilon}^2}{8\tilde{A}}.$$

The equalities $R^2_{123} = R^6_{163}$ and $R^5_{456} = R^3_{436}$ obtained from (2) lead to $\epsilon = \tilde{\epsilon} = 0$. Obviously, when $\epsilon = \tilde{\epsilon} = 0$, all the components of the curvature tensor of the h-space we consider are equal to zero.

5 The h-space of the [411] type

Metric of this space is the following

$$\begin{multline} g_{ij}dx^i dx^j = e_4\Pi_\sigma(f_\sigma - \epsilon x^4)\{6A dx^1 dx^4 + 2dx^2 dx^3 + \\ 2(2\epsilon x^2 - 3A\Sigma_1)dx^2 dx^4 - \Sigma_1 (dx^3)^2 + 2(\epsilon x^1 - 2\epsilon x^2 \Sigma_1)dx^3 dx^4 + \\ 4((\epsilon x^2)^2\Sigma_1 + \epsilon^2 x^1 x^2 - \frac{3}{2}\epsilon x^1 A\Sigma_1)(dx^4)^2\} + 3A dx^3 dx^4 + 12\epsilon x^2 A (dx^4)^2 + \sum_\sigma e_\sigma \Pi'_i (f_i - f_\sigma)(dx^\sigma)^2, \tag{18} \end{multline}$$

where

$$A = \epsilon x^3 + \theta(x^4), \quad \Sigma_1 = (f_5 - \epsilon x^4)^{-1} + (f_6 - \epsilon x^4)^{-1}, \tag{19}$$

$f_1 = f_2 = f_3 = f_4 = \epsilon x^4$, f_5 is a function of the variable x^5, f_6 is a function of the variable x^6, $e_4, e_5, e_6 = \pm 1$, ϵ equal to 0 or 1.

In this case, nonzero components of the curvature tensor have the following form: $R^i_{\nu i\mu}$ $(i \neq \mu)$, $R^{a_1}_{\nu b_1 \mu}$ $(b_1 \neq \mu, a_1 < b_1)$. Here $i = 1, \dots, 6$, $a_1, b_1 = 1, \dots, 4$, ν, μ are the indices of nonzero components of metric $g_{\nu\mu}$ (18).

Formula (2) leads to

$$R^1_{114} = R^\sigma_{1\sigma 4}, \quad R^1_{214} = R^\sigma_{2\sigma 4}, \quad R^1_{224} = 0. \tag{20}$$

For the h-space of the [411] type,

$$R^1_{114} = \rho_4 g_{14}, \quad R^1_{214} = \rho_4 g_{24}, \quad R^\sigma_{1\sigma 4} = \rho_{\sigma 4} g_{14}, \quad R^\sigma_{2\sigma 4} = \rho_{\sigma 4} g_{24},$$

$$R^1_{224} = \gamma_1 g_{24} + \gamma_2 g_{14} + \frac{2\epsilon}{3A^2}(\theta' - \frac{4}{3}\epsilon^2 x^2),$$

where

$$\gamma_1 = -\frac{1}{4}\sum_\sigma \frac{(f'_\sigma)^2}{(f_\sigma - f_4)^3 g_{\sigma\sigma}}, \quad \gamma_2 = -\frac{1}{4}\sum_\sigma \frac{(f'_\sigma)^2}{(f_\sigma - f_4)^4 g_{\sigma\sigma}}, \tag{21}$$

ρ_4 and $\rho_{\sigma 4}$ are determined by (7) when $p = 4$, $\sigma = 5, 6$. Therefore, from first two equalities of (20), we obtain

$$\rho_4 - \rho_{\sigma 4} = 0, \quad \gamma_1 g_{14} + \frac{2\epsilon^2}{3A} = 0.$$

Differentiating the last equality with respect to x^1, one obtains $\epsilon = 0$, hence, $\gamma_1 = 0$. Then, from the last equality of (20) it follows that $\gamma_2 = 0$.

Let us prove that the necessary conditions of constant curvature of the h-space of the [411] type,

$$\rho_4 - \rho_{\sigma 4} = \epsilon = \gamma_1 = \gamma_2 = 0 \tag{22}$$

are sufficient. One has

$$\partial_1 \rho_4 = \partial_2 \rho_4 = \partial_3 \rho_4 = 0,$$

$$\partial_4 \rho_4 = 6\epsilon\gamma_1, \quad \partial_\sigma \rho_4 = -f'_\sigma \frac{\rho_4 - \rho_{\sigma 4}}{f_\sigma - f_4}.$$

From these relations and taking into account (22), one obtains $\rho_4 = \rho_{\sigma 4} = \text{const}$. Using the last equality and the conditions $\epsilon = \gamma_1 = \gamma_2 = 0$, we can prove that the curvature tensor of our space satisfies identities (2) with $K = \rho_4 = \text{const}$.

6 The h-space of the $[51]$ type

Metric of the h-space of the [51] type is defined by the formulas

$$g_{ij}dx^i dx^j = e(f_6 - \epsilon x^5)\{8A dx^1 dx^5 + 2dx^2 dx^4 + \tag{23}$$

$$2(3\epsilon x^3 - 4A\Sigma_1)dx^2 dx^5 + (dx^3)^2 - 2\Sigma_1 dx^3 dx^4 + 2(2\epsilon x^2 - 3\epsilon x^3 \Sigma_1)dx^3 dx^5 +$$

$$2(\epsilon x^1 - 2\epsilon x^2 \Sigma_1)dx^4 dx^5 + 4(3/2\epsilon x^1 \epsilon x^3 + (\epsilon x^2)^2 - 2\epsilon x^1 A \Sigma_1 -$$

$$3\epsilon x^2 \epsilon x^3 \Sigma_1)(dx^5)^2\} + e_6(\epsilon x^5 - f_6)^5 (dx^6)^2,$$

where

$$A = \epsilon x^4 + \theta, \quad \Sigma_1 = (f_6 - \epsilon x^5)^{-1}, \tag{24}$$

θ is a function of the variable x^5, $f_5 = \epsilon x^5$, f_6 is a function of the variable x^6, $e_5, e_6 = \pm 1$, ϵ equal to 0 or 1.

The nonzero components of the curvature tensor have the form $R^i_{\nu i \mu}$ $(i \neq \mu)$, $R^{a_1}_{\nu b_1 \mu}$ $(b_1 \neq \mu, a_1 < b_1)$, where $i = 1, \ldots, 6$, $a_1, b_1 = 1, \ldots, 5$, ν, μ are the indices of nonzero components of metric $g_{\nu\mu}$ (23). Similarly to the previous cases, one can prove that the necessary and sufficient condition of constant curvature of the h-space of the [51] type is determined by relations

$$f'_6 = \epsilon = 0.$$

Thus, we come to the following

Theorem. *The 6-dimensional rigid h-spaces* V^6 *of the* [2211], [321], [33], [411] *and* [51] *types have the constant curvature if and only if* *for the* h*-space of the* [2211] *type*

$$\rho_p - \rho_{\sigma p} = \rho_p - \rho_{pq} = \epsilon = \tilde{\epsilon} = 0 \quad (p \neq q, p, q = 2, 4, \sigma = 5, 6), \tag{25}$$

for the h*-space of the* [321] *type*

$$f'_6 = \epsilon = \tilde{\epsilon} = 0, \tag{26}$$

for the h*-space of the* [33] *type*

$$\epsilon = \tilde{\epsilon} = 0, \tag{27}$$

for the h*-space of the* [411] *type*

$$\rho_4 - \rho_{\sigma 4} = \epsilon = \gamma_1 = \gamma_2 = 0 \quad (\sigma = 5, 6), \tag{28}$$

for the h*-space of the* [51] *type*

$$f'_6 = \epsilon = 0, \tag{29}$$

where ρ_p*,* $\rho_{\sigma p}$*,* ρ_{pq}*,* γ_1 *and* γ_2 *are determined by formulas* (7)*,* (21)*.*

I am grateful to A.V.Aminova for the constant encouragement and discussions.

References

[A] A.V. Aminova, Lie algebras of infinitesimal projective transformations of Lorentz manifolds (in Russian) // Uspekhi Mat. Nauk **50** (1995), no. 1(301), 69-142.

[E] L.P. Eizenhart, Riemannian geometry, Princeton University Press, 1997.

[P] A.Z. Petrov, On geodezic mapping of the Riemannian space with indefinite metric (in Russian)// Uch. zap. Kazan. un-ta. 1949. T. 109 no. 3. 7-36.

[Z] Z. Zakirova, PhD thesis, 2001, Kazan State University

[Z1] Z. Zakirova, First integral of the geodesics equations in 6d h-spaces // in: Proc. of Geometry Seminar, Kazan, 1996

[Z2] Z. Zakirova, 6-dimensional h-spaces of the special type // Lobachevskiy Int. geom. seminar, Kazan, 1997: Abstracts of contributed papers – Kazan, 1997.- P. 52.

[Z3] Z. Zakirova, On one solution of the Eisenhart equation. // "Latest problems of the field theory. 1999-2000". Editor A. V. Aminova.- Kazan, 2000. . 104-109.

[Z4] Z. Zakirova, First integral of the geodesics equations in h-space of the [411] type. // Izv. Vuzov. Mathematics. N9(448), 1999, . 78-79.

[Z5] Z.Zakirova, On projective group properties of the 6d h-space of the type [33], math.DG/0101052; submitted to "Classical & Quantum Gravity"